

Submitted to *Astrophys. J. Lett*, 14 April 2005;
revised 23 May 2005; accepted 27 May 2005

The γ -ray giant flare from SGR1806-20: Evidence for crustal cracking via initial timescales

Steven J. Schwartz¹

`s.schwartz@imperial.ac.uk`

and

Silvia Zane²

`sz@mssl.ucl.ac.uk`

and

Robert J. Wilson²

`rjw@mssl.ucl.ac.uk`

and

Frank P. Pijpers¹

`f.pijpers@imperial.ac.uk`

and

Daniel R. Moore³

`dan.moore@imperial.ac.uk`

and

D. O. Kataria²

`dok@mssl.ucl.ac.uk`

and

Timothy S. Horbury¹

`t.horbury@imperial.ac.uk`

and

Andrew N. Fazakerley²

anf@mssl.ucl.ac.uk

and

Peter J. Cargill¹

p.cargill@imperial.ac.uk

ABSTRACT

Soft γ -ray repeaters (SGRs) are neutron stars which emit short ($\lesssim 1$ s) and energetic ($\lesssim 10^{42}$ erg s $^{-1}$) bursts of soft γ -rays. Only 4 of them are currently known. Occasionally, SGRs have been observed to emit much more energetic “giant flares” ($\sim 10^{44} - 10^{45}$ erg s $^{-1}$). These are exceptional and rare events. We report here on serendipitous observations of the intense γ -ray flare from SGR 1806-20 that occurred on 27 December 2004. Unique data from the Cluster and Double Star-2 satellites, designed to study the Earth’s magnetosphere, provide the first observational evidence of three separate timescales within the early (first 100ms) phases of this class of events. These observations reveal that, in addition to the initial very steep (< 0.25 ms) X-ray onset, there is firstly a 4.9ms exponential rise timescale followed by a continued exponential rise in intensity on a timescale of 70ms. These three timescales are a prominent feature of current theoretical models, including the timescale (several ms) for fracture propagation in the crust of the neutron star.

Subject headings: stars: neutron – stars: individual(SGR 1806-20) – stars: magnetic fields – X-rays: stars – gamma rays: observations

¹Blackett Laboratory, Imperial College London, South Kensington, London SW7 2BW, UK

²Mullard Space Science Laboratory, University College London, Holmbury St. Mary, Dorking, Surrey RH5 6NT, UK

³Department of Mathematics, Imperial College London, South Kensington, London SW7 2BW, UK

1. Introduction

During the quiescent state (i.e., outside bursts events) Soft γ -ray Repeaters (SGRs) are detected as persistent X-ray emitters with a luminosity of $\sim 10^{35} \text{erg s}^{-1}$. Several characteristics of SGRs, including their bursting activity, are often explained in the context of the “magnetar” model (Duncan & Thompson 1992; Thompson & Duncan 1995; Thompson et al. 2002). Magnetars are neutron stars whose X-ray emission is attributed to the presence of an ultra-strong magnetic field ($\sim 10^{14} - 10^{15} \text{Gauss}$). The frequent short bursts are associated with small cracks in the neutron star crust, driven by magnetic diffusion (Thompson & Duncan 1995). Alternatively, because of their short (submillisecond) time scale, they may follow a sudden loss of magnetic equilibrium through the development of a tearing instability (Lyutikov 2002, 2003). Giant flares, acting on a much longer time scale, should involve not only the above phenomena but also global rearrangements of the magnetic field in the interior and magnetosphere of the star.

Observations made on 2004 December 27 showed that SGR 1806-20 had experienced an exceptionally powerful giant flare, lasting $\sim 0.25 \text{s}$ (Borkowsky et al. 2005; Golenetskii et al. 2004; Mazets et al. 2005; Palmer et al. 2005; Mereghetti et al. 2005). For the first 200ms it saturated almost all instruments on satellites equipped to observe γ -rays. Instruments on board the GEOTAIL spacecraft provided unique measurements (Terasawa et al. 2005) of the hard X-ray intensity throughout the event. The energy associated with γ -rays above 50 keV was $\sim 5 \times 10^{46} \text{ergs}$ (assuming a distance of 15 kpc and that it radiates isotropically), hundreds of times brighter than the giant flares previously observed from other SGRs. Following the event, a radio afterglow has been detected (Cameron & Kulkarni 2005; Gaensler et al. 2005). Several follow-on studies (e.g. Hurley et al. 2005; Israel et al. 2005) have concentrated on the characteristics of the extended tail of the flare.

In this letter we report observations of the initial flare rise and decay of SGR 1806-20. This phase was recorded by the thermal electron detectors (Johnstone et al. 1997) onboard two of the four Cluster spacecraft, and also by an identical instrument on the Chinese Double Star polar spacecraft, TC-2 (Fazakerley et al. 2005). The data reported here are unsaturated, have a slightly higher temporal resolution (4ms as opposed to 5.48ms) than that of similar measurements taken by the GEOTAIL spacecraft (Terasawa et al. 2005), and, critically, captured the initial flare activity without interruption during the very steep rise through and beyond the maximum in intensity. This dataset sheds unique information on the most intense period of flare activity.

2. Method

We have drawn on data from the PEACE instruments on Cluster-2 (C2, not shown below), Cluster-4 (C4), and Double Star Polar Orbiter (TC-2). Altogether, these spacecraft carry 5 nearly identical instruments, although the naturally occurring electrons, which the instruments are designed to detect, confuse the flare response on two of them (the Low Energy Electron Analysers on C2 and C4). As the remaining sensors provide the same coverage, with less interference, we concentrate on those. The Cluster and TC-2 spacecraft were at approximately 42,000km and 26,000km altitudes respectively and had unobstructed sight lines to the source, which was 5.3° from the Earth-Sun line (Mazets et al. 2004). Additionally, all the sensors used in the present work were not obstructed by other parts of the spacecraft during the initial flare spike.

The PEACE detector onboard TC-2 sweeps through a full range of electron energies every 121ms followed by an 8ms gap. Each sweep is divided into 30 energy channels, measured sequentially and returned in the telemetry stream, leading to a 4ms cadence in count rate values. At the time of the flare, the instrument was operating in a mode in which the lowest 14 channels were not returned, leading to a sequence of 16 measurements at 4ms resolution (64ms in total) followed by a gap of 56+8ms. The response of the instrument to X- and γ -radiation has not been calibrated, but the count rates are due to direct stimulation of the micro-channel-plate detectors at the receiving end of the electron optics. As such, it is independent of the particular energy channel and proportional to the incident flux, but dependent on the photon energy.

By contrast, the PEACE detectors onboard the other spacecraft (C2 and C4) were operating in a different mode. Firstly, neighbouring energy channels were summed pairwise, leading to an 8ms cadence within a single sweep. Moreover, neighbouring sweeps were summed pairwise. Thus an individual measurement is the sum of an 8ms sample with another 8ms sample taken 125ms later. While there is no possibility to deconvolve this data, we are able to use the steep rise and decay of the flare to our advantage by assuming that each datum is dominated by the 8ms sample located in time closest to the peak in flare intensity. The data show changes in count rate of an order of magnitude per sweep. This enables us to assign those values, with a 10% error in counts, to a unique time. However, near the peak itself, where the counting rates vary less rapidly, this technique leads to a larger error. Additionally, on Cluster the count rate due to natural electrons increases toward the end of each 125ms sweep. In this manner we reconstruct, with some unavoidable error, a time series of 8ms samples in 128ms blocks with 125ms gaps between blocks.

The response of the instruments on Cluster falls off more rapidly with time beyond the flare peak than on TC-2. This is most likely due to the changing instrumental response as

the spacecraft spins, as described below; it may also be indicative of changes in the spectral characteristics of the source (Mazets et al. 2004). We have drawn most of our conclusions from the higher resolution data on TC-2, which follows more of the event and with less natural electron interference, and used the Cluster data to fill in information about the shape of the light curve during a single 56ms period when TC-2 did not return data.

We have not attempted a determination of the absolute γ -ray flux during the event. Apart from the lack of normal ground calibrations (the instruments were not designed to measure γ -rays), the response depends on the orientation of the instrument. This is due to differences in shielding added to reduce background radiation effects on the measurements of in situ electrons, to the partial obscurations caused by other instruments, fuel tanks, and spacecraft body, to differences in the intrinsic detector response, and to the operational parameters of the detector. Some of these will vary with spacecraft spin, which has a period ~ 4 s on both C4 and TC-2. Both TC-2 and the sensor used here from C4 were relatively unobscured during the peak in intensity, but had turned through $\sim 35^\circ$ by the time of a secondary peak some 390ms later, shown below. Thus while local features are well-represented, the relative intensities of more widely separated features should be used with caution.

3. Results

Figure 1 shows the count-rate of the TC-2 electron detector over a 1.45 second interval. Zero time corresponds to the peak in count rate. We also show the previously reported GEOTAIL data (Terasawa et al. 2005), aligned in time. Where data is available from both spacecraft the detailed features match extremely well; in particular the steep rise and peak are evident. Following the first 64ms gap in the TC-2 data (see Section 2), a shoulder appeared followed by a steeper decline. Two groups of data points later, at $t \sim 390$ ms, a secondary, noisy peak can be seen (enlarged in the inset), which coincides in time and shape with that seen by the BAT instrument on SWIFT (Supplementary Figure 2 of Palmer et al. 2005) and also by GEOTAIL (Terasawa et al. 2005). Count rates return to their ambient levels ($\lesssim 10$) at a time 600ms beyond the peak in the count rate. There is no doubt that the large, but unsaturated, count rates shown in Figure 1 are signatures of the γ -ray giant flare emitted by SGR 1806-20.

There is some hint in the Cluster data (not shown) of a signal 3.5–4 seconds after the main spike. This coincides with a broad feature seen in the BAT data which marked the return of the periodic tail pulses at the same period (~ 7.56 s) as in the pre-flare observations.

In order to fill in the key time interval near the spike, we show in Figure 2 the combined data from both C4 and TC-2. The combined set reveals a dip after $t = 0$ s followed by an increase leading to the shoulder seen in the TC-2 data. Such “repeated energy-injections” on a 100ms timescale were reported in the GEOTAIL data (Terasawa et al. 2005) as can be seen in Figure 1. This 100ms timescale is a feature of all three known giant flares (Cline et al. 1980; Feroci et al. 2001). The C4 data prior to the peak show evidence of time-convolutions described in Section 2 and have been omitted from Figure 2. Accordingly, we focus mainly on the TC-2 data in what follows.

The steep initial rise is well fit by an exponential function (shown as the solid line) and has an e -folding time of 4.9ms. This is an order of magnitude longer than the 0.3ms timescales Palmer et al. (2005) find in the detailed leading edge (prior to saturation of the BAT instrument). Later, 24ms before the peak intensity was reached, the increase slowed to an exponential rise with an e -folding time of 67ms, also shown as a solid line in the figure. The previously unresolved ~ 5 ms e -folding rise marks the transition to a timescale comparable to the 100ms timescale of the overall main peak and subsequent energy injection(s).

4. Discussion and Conclusions

Thompson et al. (2002) recently proposed a scenario in which the magnetars (AXPs and SGRs) differ from standard radio pulsars since their magnetic field is globally twisted inside the star, up to a strength of about 10 times the external dipole. At intervals it can twist up the external field and propagate outward through Alfvén waves. The resulting stresses built up in the neutron star crust and the magnetic footpoint movements can lead to crustal fractures, glitching activity and flare emission (Ruderman 1991; Ruderman et al. 1998). In this scenario the giant flare is produced when the crust is unable to respond (quasi)plastically any more to the imparted stress and finally cracks (Thompson & Duncan 2001; Thompson et al. 2002); the flare emission and the crustal fracturing can be in turn accompanied by a simplification of the exterior magnetic field with a (partial) untwisting of the global magnetosphere.

We note here that all three of the timescales observed during the giant flare emitted by SGR 1806-20 can be explained in this scenario. The initial sub-ms timescale revealed by BAT is typical of reconnection processes in the external magnetosphere, where the low density gives rise to short Alfvén times. The longer ~ 100 ms timescale is related to i) the overall duration of giant flares, ii) the final part of the rise observed here and iii) the repeated intensity rises at ~ 100 ms intervals observed with GEOTAIL in the first 200ms after the peak. This time scale is comparable to the time for a 10^{15} G magnetic field to rearrange

material in the deep crust and core of the neutron star, at an internal Alfvén time, and appears as a rather regular feature in the lightcurves of giant flares (Cline et al. 1980). The intermediate ~ 5 ms time is naturally explained if the rising time is limited by propagation of a triggering fracture of size ~ 5 km given the theoretical expectation $\ell \approx 4\text{km}(t_{\text{rise}}/4\text{ms})$ (Thompson & Duncan 2001).

An alternative but extreme possibility is that the giant flare is a purely magnetospheric process in which a ~ 10 ms timescale is that which is necessary for the tearing instability to initiate large-scale magnetospheric reconnection (Lyutikov 2002, 2003). However Lyutikov (2003) concedes that the crustal fracture model is more likely to explain the extended post-flare activity. Both perspectives require an initial fast process on a magnetospheric Alfvén crossing time to explain the very steep, sub-ms rise. Thus, while the 5ms timescale reported here could be related to a tearing instability in the magnetosphere, it seems more probable that it is the first of a number of crustal events. This may not be the case for short bursts seen more frequently from SGRs, for which tearing could play an important role.

Recently, tens of Hz Quasi-Periodic Oscillations have been detected in the tail of the event (Israel et al. 2005). These modes are likely to be associated to global seismic oscillations. In particular, the large crustal fracturing inferred by us can easily excite toroidal modes with characteristic frequencies in the observed range.

The TC-2 data thus provide the clue to the missing link between the interior magnetic processes and their external consequences and probe directly the crustal properties. Cluster and Double Star were designed to study the various boundary layers of the Earth’s magnetosphere, including the physics of magnetic reconnection. Such boundary layer physics has application throughout the astrophysical plasma universe, and it is therefore appropriate that these missions contribute in a more direct way to the study of magnetic reorganisation in an astrophysical object outside the solar system.

Some of this work was supported by the UK PPARC. We are grateful to T. Terasawa for the provision of the GEOTAIL data shown in Figure 1 and for many helpful remarks. We thank G. Israel and C. Thompson for helpful comments and pre-publication material. Cluster is a project of ESA and NASA. Double Star is a project of the Chinese National Space Agency together with ESA.

REFERENCES

Borkowsky, J., et al. 2005, GRB Circular Network, 2920

- Cameron, P. B., & Kulkarni, S. 2005, GRB Circular Network, 2928, 2930, 2934, 2938, 2940, 2946, 2948
- Cline, T. L., et al. 1980, ApJ, 237, L1
- Duncan, R. C., & Thompson, C. 1992, ApJ, 392, L9
- Fazakerley, A. N., et al. 2005, Annales Geophysicae, submitted
- Feroci, M., et al. 2001, ApJ, 549,1021.
- Gaensler, B. M., et al. 2005, GRB Circular Network, 2933, 2935, 2943
- Golenetskii, S., et al. 2004, GRB Circular Network, 2923
- Hurley, K., et al. 2005, Nature, submitted
- Israel, G., et al. 2005, ApJ, submitted
- Johnstone, A. D., et al. 1997, Space Sci. Rev., 79, 351
- Lyutikov, M. 2002, ApJ, 580, L65
- . 2003, MNRAS, 346, 540
- Mazets, E., et al. 2005, GRB Circular Network, 2922
- Mazets, E. P., et al. 2004, astroph/0502541
- Mereghetti, S., et al. 2005, ApJ, submitted, astroph/0502577
- Palmer, D. M., et al. 2005, Nature, submitted, astroph/0503030
- Ruderman, M., 1991, ApJ, 382, 587.
- Ruderman, M., Zhu, T., & Chen, K., 1998, ApJ, 492, 267.
- Terasawa, T., et al. 2005, Nature, submitted, astroph/0502315
- Thompson, C., & Duncan, R. C. 1995, MNRAS, 275, 255
- . 2001, ApJ, 561, 980
- Thompson, C., Lyutikov, M., & Kulkarni, S. R. 2002, ApJ, 574, 332




Fig. 1.— (Top) Count rate returned by the PEACE instrument onboard the Chinese Double Star (TC-2) spacecraft during the intense γ -ray event of 27 December 2004. (Bottom) GEOTAIL count rates described more fully elsewhere (Terasawa et al. 2005). A steep rise, peak intensity, shoulder, and secondary peak are all evident. The signal fades into the background count-rate after ~ 600 ms. The inset shows details of the secondary peak at 390ms seen in Figure 1. This secondary peak compares well in timing and shape with the feature also seen by BAT/SWIFT (Terasawa et al. 2005; Palmer et al. 2005).




Fig. 2.— Combined Double Star (TC-2) (triangles) and Cluster-4 HEAA (gray circles) count rates. The data have been shifted in time to align features near the peak count rates. The C4 data have been reduced by a factor 2.9; only uncontaminated and weakly convoluted C4 points are shown (see Section 2). Solid lines show exponential fits to the steepest TC-2 rise, and also to the TC-2 determination of the period leading to the main peak.